

# blood

2012 120: 1367-1379  
Prepublished online June 14, 2012;  
doi:10.1182/blood-2012-05-399048

## **Allogeneic hematopoietic cell transplantation for myelofibrosis in the era of JAK inhibitors**

Vikas Gupta, Parameswaran Hari and Ronald Hoffman

---

Updated information and services can be found at:  
<http://bloodjournal.hematologylibrary.org/content/120/7/1367.full.html>

Articles on similar topics can be found in the following Blood collections

[Free Research Articles](#) (2012 articles)  
[Myeloid Neoplasia](#) (1063 articles)  
[Perspectives](#) (151 articles)  
[Transplantation](#) (1925 articles)

---

Information about reproducing this article in parts or in its entirety may be found online at:  
[http://bloodjournal.hematologylibrary.org/site/misc/rights.xhtml#repub\\_requests](http://bloodjournal.hematologylibrary.org/site/misc/rights.xhtml#repub_requests)

Information about ordering reprints may be found online at:  
<http://bloodjournal.hematologylibrary.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:  
<http://bloodjournal.hematologylibrary.org/site/subscriptions/index.xhtml>



## Perspectives

# Allogeneic hematopoietic cell transplantation for myelofibrosis in the era of JAK inhibitors

Vikas Gupta,<sup>1</sup> Parameswaran Hari,<sup>2</sup> and Ronald Hoffman<sup>3</sup>

<sup>1</sup>Princess Margaret Hospital, University of Toronto, Toronto, ON; <sup>2</sup>Medical College of Wisconsin, Milwaukee, WI; and <sup>3</sup>Mount Sinai School of Medicine, New York, NY

The discovery of *JAK2617F* mutation paved the way for the development of small molecule inhibitors of JAK1/2 resulting in first approved JAK1/2 inhibitor, ruxolitinib, for the treatment of patients with myelofibrosis (MF). Although JAK1/2 inhibitor therapy is effective in decreasing the burden of symptoms associated with splenomegaly and MF-related constitutional symptoms, it is neither curative nor effective in reducing the risk of leukemic transformation. Presently, allogeneic

hematopoietic cell transplantation (HCT) is the only curative therapy for MF. A significant risk of regimen-related toxicities, graft failure, and GVHD are major barriers to the success of HCT in MF. Because of significant HCT-associated morbidity and mortality, divergent opinions regarding its appropriate role in this clinical situation have emerged. In this review, the risk-benefit ratios of modern drug therapy compared with HCT in MF patients are analyzed. A risk-adapted ap-

proach individualized to each patient's biologic characteristics and comorbidities is described, which is currently warranted in determining optimal treatment strategies for patients with MF. Inclusion of JAK1/2 inhibitor therapy in future transplant conditioning regimens may provide an opportunity to overcome some of these barriers, resulting in greater success with HCT for MF patients. (*Blood*. 2012;120(7): 1367-1379)

## Introduction

Primary myelofibrosis (PMF) is a myeloproliferative neoplasm that originates at the level of the hematopoietic stem cell and is characterized by cytopenias, extramedullary hematopoiesis, megakaryocytic hyperplasia, reactive marrow fibrosis, and systemic symptoms resulting from elevated levels of inflammatory and proangiogenic cytokines.<sup>1</sup> The median age at diagnosis is 67 years,<sup>2</sup> and only 13% of patients are 50 years of age or younger at the time of referral.<sup>3</sup> PMF is notorious for its heterogeneity, and its clinical course varies from an indolent course persisting for almost a decade in some to others with rapidly progressive disease with a survival of 12 to 24 months.<sup>4-6</sup> A form of myelofibrosis (MF) indistinguishable from PMF can occur as part of the natural history of both polycythemia vera (PV) and essential thrombocythemia (ET), and are referred to as post-PV or ET-related MF. These 3 disorders will be collectively referred to as MF in this report.

Allogeneic hematopoietic cell transplantation (HCT) is the only curative treatment for MF at present. The therapeutic efficacy of HCT in patients with MF is mediated partly through the antineoplastic effect of pretransplant conditioning regimen and through an alloimmune GVL effect. The significant morbidity and mortality associated with HCT in MF have led to divergent opinions regarding its appropriate role.<sup>7,8</sup> Several important issues, including patient selection, timing of HCT, optimal conditioning regimens, role of prior splenectomy, and its appropriate use in older persons, remain unresolved, resulting in considerable diversity of application of HCT in MF patients.

## Evolution of HCT for MF over time

MF is a rare indication for HCT, and even major transplant centers perform limited numbers of transplantations for patients with MF. HCT was historically underused in MF as graft failure was thought

intuitively to be highly likely in the setting of marrow fibrosis. Early studies demonstrated the feasibility of engraftment as well long-term disease control in patients with MF using myeloablative conditioning (MAC) regimens.<sup>9-11</sup> The use of reduced intensity conditioning (RIC) has become widespread during the last decade. There has been a slow progressive increase in the use of HCT for MF as reflected in the trends generated by the Center for International Blood and Marrow Transplant Research (CIBMTR; Table 1; Figure 1A-B). These data provide a snapshot of current transplant practices in MF and demonstrate increasing use of HCT in older patients, greater use of peripheral blood stem cell grafts, and the growing popularity of RIC regimens.

## HCT or JAK1/2 inhibitor therapy: a changing risk-benefit ratio

Advances in supportive care, conditioning regimens, GVHD prophylaxis, and high-resolution typing for the selection of unrelated donors, have improved the safety and outcomes of HCT.<sup>12</sup> Application of RIC has expanded the scope of HCT to many older patients or those with multiple comorbidities that would have previously precluded them from the option of HCT.<sup>13</sup>

Significant progress has been made in the last few years in understanding the natural history of MF. The discovery of the *JAK2V617F* mutation in 2005 provided a significant impetus to the laboratory and translational research in MF, culminating in the Food and Drug Administration approving the first JAK 1/2 inhibitor (ruxolitinib; INCB018424) for the treatment of MF in November 2011. In a phase 3 randomized trial, 42% of patients treated with ruxolitinib experienced more than or equal to 35% reduction in spleen volume compared with 0.7% of patients receiving placebo ( $P < .001$ ) regardless of JAK2 mutational

Submitted May 15, 2012; accepted June 12, 2012. Prepublished online as *Blood* First Edition paper, June 14, 2012; DOI 10.1182/blood-2012-05-399048.

© 2012 by The American Society of Hematology

**Table 1. Time trends in HCT for myelofibrosis: data from CIBMTR**

Characteristics of patients	1997-2000, N (%)	2001-2004, N (%)	2005-2008, N (%)
No. of patients	72	213	336
No. of centers	47	114	123
<b>Age, y</b>			
≤ 9	1 (1)	0	3 (1)
10-19	1 (1)	5 (2)	8 (2)
20-29	3 (4)	3 (1)	4 (1)
30-39	12 (17)	20 (9)	11 (3)
40-49	26 (36)	76 (36)	69 (21)
50-59	25 (35)	82 (38)	151 (45)
60-69	4 (6)	25 (12)	87 (26)
≥ 70	0	2 (1)	3 (1)
<b>Donor group</b>			
HLA-identical sibling	36 (50)	111 (52)	142 (42)
Other related	3 (4)	9 (4)	12 (4)
Unrelated	33 (46)	93 (44)	182 (54)
<b>Graft type</b>			
BM	48 (67)	45 (21)	48 (14)
PBSCs	24 (33)	168 (79)	284 (85)
CB	0	0	4 (1)
<b>Conditioning regimen</b>			
Proportion of RIC	15%	37%	45%

PBSCs indicates peripheral blood stem cells; CB, cord blood; and RIC, reduced intensity conditioning.

status.<sup>14</sup> In addition, 46% of ruxolitinib-treated patients experienced a more than or equal to 50% improvement in constitutional symptoms compared with 5% in the placebo group. After a median follow-up of 51 weeks, the ruxolitinib group experienced a significant reduction in mortality (hazard ratio = 0.50; 95% CI, 0.25-0.98; *P* = .04). Patients who received ruxolitinib had mean reduction in *JAK2V617F* allele burden of 10.9% at week 24 and 21.5% at week 48; patients who received placebo had a mean

increase of 3.5% at week 24 and 6.3% at week 48. Another phase 3 trial comparing ruxolitinib with best available therapy reproduced the findings of improvement in splenomegaly and MF-related symptoms; however, a survival benefit was not demonstrated.<sup>15</sup> Two case-control studies compared the survival of patients treated with ruxolitinib with historical controls and found contradictory results.<sup>16,17</sup>

Ruxolitinib was uniformly ineffective in reversing abnormalities in peripheral blood or histopathologic abnormalities in the marrow, eliminating marker cytogenetic abnormalities or reducing the *JAK2V617F* allele burden to a degree associated with tyrosine kinase inhibitor therapy of BCR/ABL1 for chronic myeloid leukemia.<sup>14,15</sup> Other limitations have been the rapid return of splenomegaly and MF-related symptoms after discontinuation of the drug. Clinical trials with several alternative JAK inhibitors and other novel agents are underway in patients with MF to identify alternative strategies that might have more substantial effects on long-term survival.<sup>18</sup>

Because the currently available JAK inhibitor therapy is not curative, HCT remains an important therapeutic option for patients with advanced forms of MF. In this review, we evaluate the positioning of HCT in the management of MF in the light of changing risk/benefit ratios associated with HCT and emerging novel therapeutic options.


### Optimal timing of therapeutic intervention in MF

Controversy over the optimal timing of HCT for MF remains. Many patients with MF have a prolonged life expectancy and enjoy a reasonable quality of life. Therefore, exposing such patients to the immediate risk of morbidity and mortality associated with HCT is not currently thought to be appropriate. However, most patients with MF will eventually develop cytopenias, symptomatic splenomegaly, and troublesome constitutional symptoms. Some patients transform to acute myeloid leukemia (AML), which further reduces the success of HCT. Therefore, understanding the natural history of MF, and risk factors associated with leukemic transformation (LT), are vital in deciding the optimal time to consider HCT to gain the greatest benefit to the patient from such a high-risk procedure.

### Risk stratification strategies for MF

**Prognostic factors for survival.** A variety of prognostic scoring systems based on clinical characteristics have been created with the aim of identifying higher-risk patients who would benefit from HCT or experimental therapeutics. These stratification schemas were developed with the hope of minimizing treatment-related risk for patients with anticipated prolonged survival until their disease acquired characteristics associated with sufficiently shortened survival to merit the risk associated with such potentially risky therapeutic options.

Among conventional scoring systems, the Lille scoring system has been the most widely used.<sup>19</sup> Based on hemoglobin levels and the presence of leukopenia or leukocytosis, low-, intermediate-, and high-risk groups were identified with median survival of 93, 26, and 13 months, respectively. A new prognostic model known as International Prognostic Scoring System (IPSS) has been developed.<sup>4</sup> Five independent risk factors, including age more than 65 years, hemoglobin less than 10 g/dL, WBC count more than  $25 \times 10^9/L$ , peripheral blood (PB) blasts more than 1%, and



**Figure 1. CIBMTR reporting trends showing transplant activity in myelofibrosis. (A) Based on age and graft source. (B) Based on donor type.**

**Table 2. Modern risk stratification systems for survival and leukemic transformation in primary myelofibrosis**

Risk stratification system	Applicability	Prognostic factors	Risk score	Median survival, mo	Comments	
IPSS <sup>4</sup>	Diagnosis	Age > 65 y	1	Low risk (0), 135		
		Anemia (Hb < 10 g/dL)	1	Intermediate 1 risk (1), 95		
		WBC count > 25 × 10 <sup>9</sup> /L	1	Intermediate 2 risk (2), 48		
		Blood blasts > 1%	1	High (≥ 3), 27		
		Constitutional symptoms	1			
DIPSS <sup>6</sup>	Any time in disease course	Age > 65 y	1	Low risk (0), not reached	DIPSS also predicts evolution to AML, Hazard ratios of 7.8 and 24.9 for Intermediate-2 and high-risk category compared with low risk <sup>23</sup>	
		Anemia (Hb < 10 g/dL)	2	Intermediate 1 risk (1-2), 170		
		WBC count > 25 × 10 <sup>9</sup> /L	1	Intermediate 2 risk (3-4), 48		
		Blood blasts ≥ 1%	1	High (5-6), 18		
		Constitutional symptoms	1			
DIPSS plus <sup>5</sup>	Any time in disease course	DIPSS low risk	0	Low risk (0), 185	First calculate DIPSS score <sup>6</sup> and then add the score of transfusion dependency, cytogenetics, and thrombocytopenia to calculate final DIPSS plus score	
		DIPSS intermediate-1	1	Intermediate 1 risk (1), 78		
		DIPSS intermediate-2	2	Intermediate 2 risk (2-3), 35		
		DIPSS high-risk	3	High (≥ 4), 16		Based on single institution data, needs validation
		<i>PLUS</i>				
		Transfusion dependency	1			
		Unfavorable cytogenetics*	1			
		Platelets < 100 × 10 <sup>9</sup> /L	1			
IWG risk model for LT <sup>25</sup>	Any time in disease course	Risk factors: high-risk cytogenetics†	2	Risk of LT: low (score 0), 3% at 3 y		
		Risk factors: blood blasts ≥ 2%	1	Risk of LT: intermediate (score 1), 10% at 3 y		
		Risk factors: platelets ≤ 50 × 10 <sup>9</sup> /L	1	Risk of LT: high (score 2), 35% at 3 y		

Hb indicates hemoglobin; WBC, white blood cell; and IWG, International Working Group.  
 \*Unfavorable cytogenetics includes +8, -7/7q-, i(17q), -5/5q-, 12p-, Inv(3) or 11q23.  
 †High-risk cytogenetics was defined as monosomal karyotype, Inv 3 or i(17q).

presence of constitutional symptoms at the time of diagnosis were predictive of survival of patients with PMF at the time of diagnosis. The presence of 0, 1, 2, and more than or equal to 3 factors are categorized as low-, intermediate-1-, intermediate-2-, and high-risk disease with a median survival of 135, 95, 48, and 27 months, respectively (Table 2).

The risk factors of IPSS were also analyzed in a time-dependent fashion termed dynamic IPSS (DIPSS).<sup>6</sup> Acquisition of anemia had a higher adverse impact on survival (roughly double) compared with other factors; therefore, anemia was assigned a score of 2 (Table 2). Thus, DIPSS differs from IPSS, which gave the same weight to each risk factor.

Marked interpatient variability within IPSS and DIPSS risk groups is observed, suggesting a potential role of other risk factors for precise risk stratification. Cytogenetics, transfusion dependency, and thrombocytopenia were incorporated into the DIPSS plus scoring system (Table 2).<sup>5</sup> DIPSS plus has not been to date validated in another independent dataset.

For risk stratification, it is recommended to use IPSS at the time of diagnosis and the DIPSS anytime during the disease course in patients with PMF.<sup>6</sup>

**Risk of LT.** LT after MF results in a resistant form of AML that is almost universally associated with poor outcomes.<sup>20-22</sup> One of the goals of HCT for PMF patients is to transplant them before LT to avoid this complication. Patients with intermediate-2 and high-risk DIPSS groups had a 7.8-fold and 24.9-fold risk of LT compared

with patients with low-risk disease.<sup>23</sup> Patients with either thrombocytopenia (platelet count < 100 × 10<sup>9</sup>/L) or an unfavorable karyotype (as discovered in “Cytogenetics”) had 18% and 31% risk of LT at 5 years and 10 years, respectively.<sup>5</sup> The corresponding risk of LT was 6% and 12% in patients with platelet count more than or equal to 100 × 10<sup>9</sup>/L and not having unfavorable karyotype. It also appears that risk of LT may be higher in patients with transfusion dependency.<sup>24</sup> The risk of LT has been further evaluated by introducing a weighted scoring system.<sup>25</sup> This proposed system described 3 risk factors for LT: high-risk karyotype [defined as monosomal karyotype, Inv 3 or i(17q); score 2]; PB blasts more than or equal to 2% (score 1); and thrombocytopenia (platelet count ≤ 50 × 10<sup>9</sup>/L; score 1).<sup>25</sup> Based on scores of these 3 factors, there were 3 distinct groups with scores of 0, 1, and more than or equal to 2, and corresponding risk of LT at 3 years was 3%, 10%, and 35%, respectively.

Available data highlight the importance of cytogenetics, PB blasts, and severity of thrombocytopenia in predicting LT in PMF.

**Risk stratification in post-PV MF and post-ET MF.** It is important to note that IPSS, DIPSS, and DIPSS plus have been studied in patients with PMF. These scoring systems have not been validated in patients with post-PV MF and ET-related MF. Available studies addressing this topic are limited by small sample size.<sup>26,27</sup> At present, same scoring systems are being used for risk stratification in patients with PPV-MF and PET-MF. Efforts are in progress to develop prognostic models for these patients. Until

**Table 3. Impact of cytogenetics and molecular prognostic markers in patients with myelofibrosis**

Cytogenetics/molecular markers	Frequency, %	Impact on LT	Impact on survival	Comments
Diploid karyotype <sup>32,33</sup>	57-71	7% at 5 y	46 mo	
<b>Sole abnormalities, standard risk</b>	15-22	Comparable with diploid karyotype	Comparable survival with diploid karyotype	
Sole del 20q <sup>32,33</sup>	4-7			Association noted between sole 20q- and leucopenia and thrombocytopenia <sup>32</sup>
Sole del 13q <sup>32,33</sup>	2-4			
Sole +9 <sup>32,33</sup>	2-3			
Chromosome 1 translocation/duplication <sup>32</sup>	2-3			
Other miscellaneous sole abnormalities other than listed under unfavorable risk <sup>32</sup>	5			
<b>Sole abnormalities, unfavorable risk</b>	4-7	Significant high risk of LT compared with diploid karyotype (46% vs 7% at 5 y)	Poor survival comparable to diploid karyotype (15 mo vs 46 mo)	
Sole +8 <sup>30,32</sup>	2-3			
Sole -7/7q-/-5/5q-/Inv 3 <sup>32</sup>	1-2			
Chromosome 17 abnormalities <sup>33</sup>	1-2			
<b>Double abnormalities*</b>				
Standard risk <sup>32</sup>	2-3	Comparable with diploid karyotype	Similar to diploid karyotype	
		Significant high risk of LT compared with diploid karyotype (46% vs 7% at 5 y)	Similar to complex karyotype	
<b>Double abnormalities†</b>				
Unfavorable risk <sup>32</sup>	2-3			
<b>Complex karyotype (≥ 3 abnormalities)<sup>32,33</sup></b>	6-8	Significant high risk of LT compared with diploid karyotype (46% vs 7% at 5 y)	Poor survival comparable to diploid karyotype (15 mo vs 46 mo)	~ 50% patients with CK have monosomal karyotype with very poor survival <sup>30</sup>
JAK2V617F <sup>41-44</sup>	55-60	No difference compared with WT	No difference compared with WT	Low JAK2V617F allele burden associated with inferior survival <sup>42,43</sup> ; and associated with myelodepletive type of MF
MPL <sup>45</sup>	8-10	No difference compared with JAK2V617F mutation positive or unmutated JAK2/MPL	No difference compared with JAK2V617F mutation positive or unmutated JAK2/MPL	MPL and JAK2 unmutated patients were significantly younger than JAK2-mutated patients <sup>45</sup>
LNK <sup>39</sup>	2-3	Not known	Not known	
TET2 <sup>66</sup>	7-17	No	No	More frequent in older patients compared with younger patients <sup>66</sup>
DNMT3A <sup>87</sup>	10-15	No	No	
IDH <sup>88</sup>	4	Shortened LFS in PMF	Short survival in PMF	
EZH2 <sup>69</sup>	6-9	Shorter LFS in patients with PMF with EZH2 mutations	Associated with short survival in idiopathic MF, independent of IPSS	Associated with higher leukocyte count, blast cell counts, and larger splenomegaly
SUZ12/EED <sup>90,91</sup>	1-3	Not clear	Not clear	May be associated with LT

\*Two abnormalities, excluding unfavorable.

†Two abnormalities, including unfavorable.

more data become available, risk stratification models used for PMF may be used for enrollment of these patients in clinical trials and therapeutic decision making.

Current risk models for survival and LT have been generated from retrospective studies in an era when therapeutic options were limited and HCT rarely used. Prospective validation and further improvement on these models by including other important variables, such as comorbidities in the setting of improving treatment options, will be desirable in future.

### Cytogenetics and molecular aberrations in MF

**Cytogenetics: a growing appreciation of its utility for prognostication of MF patients.** Several reports have highlighted the impact of cytogenetic abnormalities on the outcomes of patients with

MF.<sup>28-31</sup> Commonly observed cytogenetic abnormalities in MF and their prognostic impact are summarized in Table 3. Cytogenetic abnormalities are seen in approximately 35% to 43% of patients with MF.<sup>31-33</sup> Karyotypic abnormalities involving one chromosome (sole), 2 chromosomes (double), or more than or equal to 3 chromosomes (complex) are observed in approximately 70%, 15%, and 15% patients, respectively.<sup>32</sup> Based on a large series of 433 patients with PMF, a 2-tiered cytogenetic risk stratification system has been proposed identifying favorable and unfavorable karyotypes.<sup>32</sup> Favorable karyotypes include: normal, sole 20q-, sole 13q-, sole chromosome 1 translocation/duplication, sole +9, other sole abnormalities (excluding those with unfavorable risk), and 2 abnormalities excluding unfavorable ones. Unfavorable karyotypes include: complex (≥ 3 abnormalities), sole +8, sole -7/7q-, sole 5/5q-,

i(17q), inv(3), 12p- or 11q23, and 2 abnormalities including an unfavorable type. Patients with favorable and unfavorable karyotypes have median survivals of 5.2 years and 2 years, respectively and corresponding risk of LT at 5 years is 7% and 46%, respectively.

Our understanding of cytogenetics in MF is at early stages compared with other myeloid malignancies, and the scheme described in previous paragraph is likely to be validated and refined in the future.

### Molecular aberrations in MF: are they useful in prognostication?

The majority of patients with MF have hematopoietic cells that are characterized by overactivation of the JAK-STAT pathway or mutations affecting chromatin structure (Table 3). *JAK2 V617F* is the most common mutation observed in MF patients.<sup>34-37</sup> Additional mutations include *JAK2 exon 12*, *MPL*, and *LNK*.<sup>38-40</sup> The prognostic significance of *JAK2V617F* has been evaluated in several studies.<sup>41-44</sup> One study described the adverse impact of *JAK2V617F* in patients with PMF,<sup>41</sup> and another study described a higher rate of LT in patients with *JAK2V617F*.<sup>44</sup> Other studies with larger sample sizes have not shown a significant difference between *JAK2* mutated and unmutated patients.<sup>42,43</sup> Importantly, 2 studies have demonstrated the shortened survival of *JAK2V617F*-positive patients with a low allele burden,<sup>42,43</sup> indicating that a low *JAK2V617F* allele burden was associated with a myelodepletive variant of PMF. Overall, the prognostic significance of *JAK2V617F* mutation remains unclear. The presence of *MPL* mutations does not impact survival or LT in PMF.<sup>45</sup>

As highlighted in Table 3, the biologic consequences of most of the other known mutations remain unclear. The oncogenetic events that transform myeloproliferative neoplasm (MPN) to AML are poorly characterized. Several genes were implicated in LT, as evidenced by mutational analysis of 63 patients with AML secondary to a preexisting MPN.<sup>46</sup> Frequent mutations were identified in *TET2* (26.3%), *ASXL1* (19.3%), *IDH1* (9.5%), and *JAK2V617F* (36.8%) mutations in AML, and all possible mutational combinations of these genes were observed. Analysis of 14 patients with paired samples during chronic-phase MPN and subsequent AML revealed that *TET2* mutations were frequently acquired at the time of LT (6 of 14; 43%). In contrast, *ASXL1* mutations were almost always detected in both the MPN and AML clones from individual patients. Mutations in *TET2*, *ASXL1*, and *IDH1* were common in MPN-related AML. Although *TET2/ASXL1* mutations may precede acquisition of *JAK2* mutations by the MPN clone, mutations in *TET2*, but not *ASXL1*, are commonly acquired at the time of LT. These findings indicate that the mutational order of events in MPN and sAML varies in different patients and that *TET2* and *ASXL1* mutations have distinct roles in MPN pathogenesis and LT. Because some cases of AML have no preexisting *JAK2/TET2/ASXL1/IDH1* mutations, it is probable that there are other mutations that are necessary for LT. Recently, recurrent mutations in the serine/arginine-rich splicing factor 2 (*SRSF2*) gene were described in AML transformed from MPNs.<sup>47</sup> At present, none of these mutations can be used in developing more robust risk stratification for either survival or risk of LT.

Identifying higher-risk patients will aid in more accurate decision making. Every patient with MF should have a detailed risk assessment at regular intervals on an ongoing basis using modern risk stratification systems (Table 2).

## Nontransplant therapeutic options in MF

### Conventional options: limited efficacy and scanty prospective evaluation

Clinical symptoms of patients with MF can be grouped into 3 main categories: cytopenias (mainly anemia), splenomegaly, and constitutional symptoms. Nontransplant treatment options mainly include supportive therapy and the use of various drugs for symptomatic improvement. Hydroxyurea (HU) is the most commonly used conventional treatment for patients with MF and is sometimes useful in managing some of the hyperproliferative manifestations of MF, such as splenomegaly, extreme leukocytosis, and thrombocytosis.<sup>48</sup> Clinical improvement with HU according to International Working Group for Myeloproliferative Neoplasms Research and Treatment criteria was seen in 40% of patients, and median duration of response was 13.2 months. Anemia or new-onset pancytopenia was observed in 45% patients. The usefulness of HU has been recently questioned. HU was the most commonly used best available therapy in COMFORT-II trial, and none of the patient treated with HU met the primary endpoint of more than or equal to 35% reduction in spleen volume.<sup>15</sup> Various other agents, such as erythropoiesis-stimulating agents, androgens, busulfan, anagrelide, interferon, and corticosteroids, and immunomodulatory derivatives, such as thalidomide or lenalidomide, have been used with mixed success in MF.<sup>8</sup> Limited efficacy and scanty prospective clinical data have prevented defining the exact role of each of these agents in the management of MF. None has been shown to modify the natural history of the disease, and their use is mainly physician dependent.

Other treatments for advanced splenomegaly include splenectomy and low-dose radiation therapy. Historically, splenectomy has been performed in approximately 10% of patients with MF in the pre-JAK inhibitor era<sup>4</sup> and is associated with significant risk of perioperative complications (27.7%) and mortality (6.7%).<sup>49</sup> With the wider availability of JAK inhibitors, the option of splenectomy is likely to decrease and will be used in selected patients who are unable to tolerate JAK inhibitor therapy because of severe cytopenias. Splenic radiation results in temporary short responses and is associated with significant cytopenias.

### Novel drugs in MF

***JAK1/2 inhibitor therapy: benefits and limitations.*** The clinical benefits of the JAK1/2 inhibitors are related to reducing the burden of troublesome symptoms of MF by reduction in splenomegaly and amelioration of constitutional symptoms.<sup>14,15,50-52</sup> These agents mainly inhibit dysregulated JAK-STAT signaling present in *JAK2V617F*-positive and -negative patients and therefore are equally efficacious irrespective of JAK2 mutation status. Anti-JAK1-mediated reduction of proinflammatory and proangiogenic cytokines is an important effect of these drugs.<sup>52,53</sup> However, there are limited effects on survival, resolution of marrow fibrosis, cytogenetic abnormalities, *JAK2V617F* allele burden, or LT, indicating a lack of effect on disease progression. Other limitations of current JAK1/2 inhibitors are the occasional return of MF-related symptoms on their discontinuation, unpredictable response duration, and lack of long-term safety and efficacy data. Nevertheless, JAK inhibitor therapy is an important advance for patients with MF and has significant clinical value in decreasing the symptom burden and improvement in quality of life.

Apart from JAK1/2 inhibitors, several other novel agents are at various stages of clinical development in MF. The most prominent among these are third-generation immunomodulatory derivative pomalidomide (phase 3), histone deacetylase inhibitors, such as panobinostat and givonostat (phase 2), mTOR inhibitors (phase 1 or 2), inhibitor of hedgehog pathway (Saridegib, phase 1), AB0024, a monoclonal antibody inhibiting LOXL2 (phase 1), and a TGF- $\beta$  signaling inhibitor (phase 1).

---

## Transplantation for MF

Major studies of HCT outcomes with more than 20 patients with MF in chronic phase, and published in peer-reviewed journals are summarized in Table 4. These studies, except one,<sup>54</sup> are retrospective in nature. Another prospective study completed by Myeloproliferative Diseases-Research Consortium has not been published in a peer-reviewed journal yet.<sup>55</sup>

### The early era of HCT in MF: feasible and curative, but associated with a high mortality

In the early era of HCT for MF, outcomes of relatively small number of patients were reported.<sup>9-11</sup> These studies established the feasibility and curative potential of HCT in MF and demonstrated that severe marrow fibrosis was not a barrier to engraftment.<sup>9-11</sup> High-dose cyclophosphamide with busulfan or total body irradiation were the most commonly used conditioning regimens.<sup>56</sup> Several important observations were made from these studies. Splenomegaly and marrow fibrosis resolved slowly in the majority after successful engraftment. Regimen-related toxicities and nonrelapse mortality (NRM) were high, and expected long-term survival was in the range of 30% to 40%, restricting the use of this option to younger patients. Patients older than 50 years, those receiving alternative donor grafts, and those with higher-risk MF were more likely to experience treatment failure. The outcomes from these reports performed more than 15 to 20 years ago are not generally applicable today, although they are sometimes used as justification for the reluctance to refer MF patients for transplantation.

### The modern era of HCT: novel conditioning strategies and establishing the feasibility of reduced intensity conditioning

During the last decade, several important advances have been made in the traditional MAC regimens. Advances, such as the use of intravenous busulfan, targeted dose of busulfan, and reversed order conditioning with cyclophosphamide followed by busulfan, reduced early regimen-related toxicities associated with HCT.<sup>10,12,57,58</sup>

The introduction of a variety of newer and lower-intensity conditioning regimens shifted the emphasis of pretransplantation conditioning therapy from myeloablation to establishing an immune-suppressive effect sufficient to establish donor hematopoiesis. These regimens, developed with the aim of reducing transplantation-related morbidity and mortality, were rapidly adopted in MF. Several lines of data suggest an immunologically mediated GVL effect as the reported success of donor-lymphocyte infusions in MF, and the success of RIC in establishing donor engraftment and resolution of marrow fibrosis over time.<sup>54,59-61</sup>

Even among the lower-intensity regimens, intensity can vary from minimal (usually 20%-25% of full intensity) or truly nonmyeloablative to reduced intensity (~40%-50% of full intensity).<sup>62</sup> We will use the term RIC collectively for minimal intensity and reduced intensity regimens here. Published literature and CIBMTR

reporting trends indicate that fludarabine in combination with busulphan/melphalan or total body irradiation are the most commonly used RIC regimens in patients with MF.<sup>54,55,63-66</sup> At present, there are no data to indicate the superiority of one regimen over the other.

### Intensity of conditioning therapy for MF: is RIC better than MAC?

There are no prospective studies comparing MAC and RIC in MF. Several retrospective studies have compared MAC and RIC regimens.<sup>56,64,65,67</sup> Similar outcomes were reported in all except one.<sup>67</sup> This study reported a more favorable outcome of patients younger than 60 years undergoing RIC transplantation.<sup>67</sup> Lack of statistical power, retrospective nature, and long time interval studied are major issues in these reports. Moreover, there are significant differences in patient populations as patients undergoing RIC are usually much older, have significant comorbidities, and have worse performance scores. NRM with RIC regimens are usually in the range of 15% to 20%; however, relapse is a major cause of failure observed in approximately 30% to 35% patients (Table 4).

A snapshot of various recently reported studies would indicate that progression-free survival is observed in approximately 45% to 50% at 3 years in patients undergoing HCT using modern conditioning regimens (Table 4).

---

## Prognostic factors for outcome of HCT in MF

The outcome of HCT is usually determined by a complex summation of various patient-, disease-, and transplant-related factors. Data validating the utility of prognostic factors in predicting the outcome of MF patients undergoing HCT are limited and conflicting.

### Patient-related factors

Age has been identified as an important prognostic factor for survival in several transplantation studies.<sup>11,54,58,67-69</sup> In the early transplantation era, high NRM was a barrier for successful outcome in patients older than 50 years. With modern conditioning regimens and RIC, many centers consider the option of HCT in the sixth and seventh decade.<sup>13</sup> Age more than 55 years/more than or equal to 57 years was identified as an independent prognostic factor (overall mortality 2.7 times) in studies from Germany using RIC consisting of fludarabine and busulfan.<sup>54,68</sup> Performance status at HCT<sup>56</sup> and high comorbidity scores<sup>58</sup> are other important patient-related factors affecting outcomes of HCT. The burden of comorbidities is usually higher in older patients.<sup>57</sup> Therefore, it is prudent that older patients are selected for HCT only after careful evaluation of performance status and comorbidities.

### MF-related factors

***Do the independent risk factors described in IPSS/DIPSS/DIPSS plus scores have similar prognostic value for patients undergoing HCT for MF?*** Anemia (hemoglobin < 100 g/dL)<sup>11,57</sup> and a greater number of transfusions before HCT (> 20 U)<sup>63</sup> have been identified as predictors for inferior survival after HCT. However, these findings have not been confirmed in other studies.<sup>68</sup> WBC count, an important independent prognostic marker in all the risk

**Table 4. Summary of major reports on HCT outcomes in MF**

Reference	Timeline of HCT	N	Median age, y (range)	Conditioning regimen	% of patients with RIC	% with MRD	NRM	PFS	OS	Comment
Guardiola <sup>11</sup>	1979-1997	55	42 (4-53)	TBI based (63%)	0	90	27% at 1 y	39% at 5 y	47% at 5 y	Hb < 10 g/dL and osteomyelofibrosis associated with lower survival
Deeg <sup>10</sup>	1980-2002	56	43 (10-66)	Bu/Cy in 78%	0	64	14% at 3 mo	NR	58 at 3 y	Targeted Bu use improved survival; cGVHD 59% at 2 y
Daly <sup>9</sup>	1990-2002	25	48 (45-50)	TBI based (92%)	0	52	48% at 1 y	NR	41 at 2 y	Prohibitive NRM; no benefit of splenectomy
Rondelli <sup>66</sup>	NR	21	54 (27-68)	Multiple	100	85	10% at 1 y	81% at 2.7 y	85% at 2.7 y	Extensive cGVHD in 44%; 2 patients needed DLI for 100% donor chimerism; resolution of fibrosis and splenomegaly in majority
Kerbauf <sup>69</sup>	NR	104	49 (18-70)	Multiple, Bu/Cy (62%)	9	50	35% at 5 y	NR	61% at 5 y	3 syngeneic donors, 54 of the patients overlapped with a prior report <sup>10</sup> ; targeted Bu improved OS; comorbidity score had impact on survival
Patriarca <sup>65</sup>	1986-2006	100	49 (21-68)	Multiple, Bu/Cy 50% of full intensity; Thiotepa + Cy in 46% of RIC	52	78	43% at 3 y	35% at 3 y	42% at 3 y	AHCT before 1995; unrelated donor and longer interval from diagnosis predicted worse outcome but not conditioning intensity; relapse at 2 y 41%, progressive decline in NRM over 20 y studied
Kroger <sup>64</sup>	2002-2007	103	55 (32-68)	Flu-Bu (100%)	100	32	16% at 1 y	51% at 5 y	67% at 5 y	First prospective study in MF, cGVHD in 43%; 12% NRM for fully matched donor AHCT; age > 55 y and HLA mismatch adversely affected OS; JAK2-positive recipients had better EFS and OS; splenectomy increased risk of relapse
Gupta <sup>64</sup>	1998-2005	46	47 y MAC; 54 y RIC	Multiple, Cy TBI (96%) for MAC; Flu Bu (70%) for RIC	50	54	48% for MAC and 27% for RIC at 3 y	43% for MAC and 58% for RIC at 3 y	48% for MAC and 68% for RIC at 3 y	RIC recipients had more advanced disease and poor KPS; low risk of relapse after either conditioning; lower GVHD with novel conditioning possibly related to use of ATG
Ballen <sup>56</sup>	1989-2002	289	47 (18-73)	Multiple, Bu/Cy (43%)	21	56	35% siblings 50% for URD at 5 y	33% siblings 27% for URD at 5 y	37% siblings 30% for URD at 5 y	Relapse at 5 y, 32% in sibling and 23% in URD; performance status, peripheral blasts sibling donor status impacted OS; RIC was similar in outcomes, except early NRM
Alchalby <sup>71</sup>	1999-2009	162	56 (32-73)	Flu-Bu in 96%	100	27	22% at 1 y	46% at 5 y	62% at 5 y	82 patients reported previously <sup>64</sup> ; age and HLA mismatch impacted NRM; 23% relapse at 3 y; clearance of mutated JAK2 at median of 96 days, and this reduced relapse risk
Bacigalupo <sup>63</sup>	1994-2007	46	51 (24-67)	Thiotepa-Cy + melphalan	100	65	24% at 5 y	NR	45% at 5 y	A risk score based on transfusion history, spleen > 22 cm and alternative donor use predicted lower OS; no benefit for splenectomy
Stewart <sup>76</sup>	1989-2005	51	49 (19-64)	Multiple, RIC in 47%	47	65	41% at 2 y	44% and 24% at 3 y for MAC and RIC	44% and 31% at 3 y for MAC and RIC	
Robin <sup>82</sup>	1997-2008	147	53 (20-68)	Multiple	69	61	39% at 4 y	32% at 4 y	39% at 4 y	19% patients had LT; poor outcome with mismatched donor
Samuelson <sup>13</sup>	1999-2007	30	65 (60-78)	Multiple	63	50	13% at day 100	40% at 3 y	45% at 3 y	Studied outcomes in patients ≥ 60 y, 7 patients had preceding LT
Abelsson <sup>67</sup>	1982-2009	92	46 for MAC, 55 for RIC	Multiple	56	40	32% for MAC and 24% for RIC at 2 y	NR	49% for MAC and 59% for RIC at 5 y	Overall NRM similar, but 5-year OS superior for RIC younger than age 60 years; less advanced MF associated with better OS
Nivison-Smith <sup>63</sup>	1993-2005	57	47 (16-71)	Multiple	26	68	25% at 1 y		58% at 5 y	Poor outcome in patients > 50 y
Ditschkowski <sup>70</sup>	1994-2010	76	50.5 (22-67)	Multiple	NR	35	36% at 5 y	50% at 5 y	53% at 5 y	Significant high risk of relapse in patients without cGVHD; DIPSS was predictive of survival
Scott <sup>57</sup>	1990-2009	170	51.5 (12-78)	Multiple	NR	50	34% at 5 y	57% at 5 y	57% at 5 y	Post-HCT success was dependent on pre-HCT DIPSS scores

MRD indicates matched related donor; PFS, progression-free survival; OS, overall survival; TBI, total body irradiation; Hb, hemoglobin; Bu, busulfan; Cy, cyclophosphamide; cGVHD, chronic GVHD; NR, not reported; DLI, donor lymphocyte infusion; AHCT, allogeneic HCT; EFS, event-free survival; KPS, Karnofsky performance status; and ATG, antithymocyte globulin.



stratification systems, did not impact transplant outcomes.<sup>57,68</sup> The presence of blasts in PB was associated with inferior outcomes,<sup>56,57</sup> mainly because of higher risk of relapse.<sup>57,71</sup> However, prognostic value of PB blasts was not confirmed in a large study of patients treated with RIC.<sup>68</sup> The presence of constitutional symptoms before HCT was associated with 2.8-fold higher risk of mortality in one study<sup>68</sup> but did not influence the outcomes in another study.<sup>57</sup> Adverse impact of cytogenetics in MF patients, described by some,<sup>10,11,70</sup> has not been confirmed by others.<sup>58,65,68</sup> Patients with abnormal cytogenetics are a small proportion in these studies for making meaningful conclusions. Thrombocytopenia ( $< 100 \times 10^9/L$ ) also appears to be an independent poor prognostic factor for survival in HCT recipients.<sup>58</sup>

**Do other MF-related factors have any prognostic value in HCT patients?** The impact of *JAK2V617F* mutation status on the outcome of transplantation is not clear. Higher overall mortality and increased risk of graft dysfunction observed by some investigators in *JAK2* wild-type patients<sup>68,72</sup> were not confirmed by other groups.<sup>70-72</sup> *JAK2* mutational status may be used as a marker of minimal residual disease. Patients who were still positive for *JAK2V617F* mutation at 6 months after HCT had a significantly greater risk of progression (5% vs 30%).<sup>71</sup>

Higher grades of fibrosis have been associated with poor outcomes in the earlier studies,<sup>11,56</sup> although the independent value of this variable has not been established.<sup>10,11,56,58,68</sup> These studies are further limited by the lack of a uniform assessment of grading of fibrosis. Longer intervals between diagnosis and HCT have been associated with mortality in one study<sup>65</sup>; however, they were not confirmed in a further studies.<sup>57,68</sup> Splenomegaly more than 22 cm has been associated with poor survival.<sup>63</sup>

### Prognostic value of different risk stratification systems in HCT patients

The prognostic value of the Lille scoring system has been most extensively studied in HCT recipients.<sup>10,54,65,67,73</sup> The patients with low-risk disease have better outcomes compared with intermediate- and high-risk patients. There appears to be a higher risk of relapse with high Lille scores.<sup>54</sup>

The Seattle group recently evaluated the usefulness of DIPSS score in HCT recipients.<sup>57</sup> The HRs of post-HCT mortality and NRM were 4.11 and 3.41 among DIPSS high-risk patients compared with low-risk patients. Median survival was not reached for DIPSS low and intermediate-1 risk cohorts, whereas it was 7 and 2.5 years, respectively, for the intermediate-2 and high-risk groups. This study highlights the dilemma of transplantation in MF in that those transplanted earlier in the course of are most likely to be cured, whereas these same patients are the ones who least need HCT, as the risk of death or LT from MF is low.<sup>5,24</sup>

A German group recently compared various risk models in a patients treated with a uniform RIC regimen.<sup>68</sup> Advanced age ( $> 57$  years), *JAK2* V617F wild-type status, and constitutional symptoms were predictive of poor survival. For those with all 3 risk factors, the hazard of death was increased by 16-fold. The DIPSS system, although predictive, did not sufficiently distinguish between intermediate-1 and intermediate-2 risk groups in this study.

### Transplant-related factors

Similar outcomes have been reported in patients undergoing matched sibling donor (MSD) and well matched (10 of 10) unrelated donor (URD) transplantation.<sup>54,58</sup> The outcomes of mismatched donors are significantly inferior.<sup>54</sup> Haploidentical or cord blood

grafts are important alternative graft sources for hematologic malignancies. However, their utility has not been well established in patients with MF. A small study described the use of cord blood grafts in patients with severe marrow fibrosis; however, 11 of 14 patients in this study had fibrosis associated with AML or myelodysplastic syndrome,<sup>74</sup> which are biologically distinct diseases than MF. There does not appear to be significant difference in outcomes of peripheral blood stem cell and marrow grafts in MF patients.<sup>56,57</sup>

The data herein highlight the difficulties in interpreting the conflicting results between various studies and making informed decisions. The reasons for such conflicting data are retrospective nature of studies, heterogeneity among patients, small sample sizes lacking statistical power, and thus inability to analyze these factors in multivariate analysis.

### Splenectomy before HCT in MF: should we or should we not?

Conflicting data exist on the effect of pre-HCT splenectomy on relapse and survival. Earlier studies from several groups did not show an impact of prior splenectomy on survival<sup>9,56,58,75,76</sup>; however, recent studies evaluating a larger number of patients demonstrated beneficial effect of splenectomy on survival.<sup>57,77</sup> Relapse risk was higher in splenectomized patients in a German study,<sup>54</sup> whereas no impact was observed in the CIBMTR study.<sup>56</sup> The discrepant results in various studies may be related to selection biases, as fitter patients who are able to undergo successful splenectomy may have a lower mortality after HCT. In addition, patients who have larger spleens may have more advanced disease, which may explain higher relapse rates.<sup>54</sup> A consistent finding reported in several studies is the faster hematopoietic recovery in splenectomized patients.<sup>11,58,76,78</sup> The procedure of splenectomy is associated with significant risk of perioperative complications (27.7%) and mortality (6.7%).<sup>49</sup> Another disadvantage of splenectomy is immunologic issues in the post-HCT setting (eg, poor response to vaccines). Reduction of splenomegaly by *JAK1/2* inhibitor therapy before HCT may be a reasonable alternative to splenectomy but without the risks of surgery related morbidity and mortality.

Given the lack of favorable data and significant perioperative complications associated with this procedure, routine splenectomy is not recommended before HCT.

---

## Barriers to success of HCT in MF

### Regimen-related toxicities: higher risk of early hepatotoxicity

Patients with MF are at a significantly higher risk of developing early hepatotoxicity.<sup>79</sup> A case-control study from Toronto evaluated early hepatotoxicity in 53 patients undergoing HCT for MF.<sup>79</sup> Compared with matched myelodysplastic syndrome patients, patients with MF had a significantly higher risk of moderate/severe hyperbilirubinemia (44% vs 21%,  $P = .02$ ) and veno-occlusive disease (36% vs 19%,  $P = .05$ ). Moderate/severe hyperbilirubinemia had an adverse impact on survival. Investigators hypothesized that a higher rate of hepatotoxicity in MF patients may be related to underlying asymptomatic portal hypertension. Screening for asymptomatic portal hypertension, using upper gastrointestinal endoscopy and abdominal Doppler ultrasound, should be considered in the pretransplantation workup of patients with MF.

### Graft failure

The incidence of GF in MF patients is reported between 5% and 25%.<sup>55,56,64</sup> In a large CIBMTR study, GF was significantly higher in patients undergoing HCT using donors other than MSD (MSD, 9%; other related, 27%; and unrelated, 20%).<sup>56</sup> Similar trends were observed in a prospective study from Myeloproliferative Diseases-Research Consortium, which showed a significantly higher rate of primary GF in MF patients undergoing URD transplantation compared with MSD (25% vs 3%).<sup>55</sup> TNF- $\alpha$  is a negative regulator of expansion and renewal of normal hematopoietic stem cells<sup>80,81</sup> and may have a differential effect on MPN and normal hematopoietic cells.<sup>82</sup> Patients with advanced MF have increased plasma levels of TNF- $\alpha$ .<sup>83</sup> It is possible that the higher rate of GF may be related to cytokines associated with more advanced disease rather than donor factors alone. Biology and risk factors for GF are poorly understood in MF and need to be studied in well-designed studies.

### GVHD

GVHD is one of the most debilitating complications of HCT. In the published literature, some studies have reported higher than expected rates of acute GVHD in MF patients,<sup>10,11,55,58,64</sup> whereas in others the rates of GVHD do not appear much different from other hematologic malignancies.<sup>54,56</sup> The impact of conditioning on acute GVHD is not clear in MF. In a prospective study from Myeloproliferative Diseases-Research Consortium, the incidence of severe acute GVHD (grade 3 or 4) was 12% and 21% in patients undergoing RIC transplantation using MSD and URD, respectively.<sup>55</sup> An alarmingly high incidence of severe GVHD in URD transplants in this study is of concern despite the use of thymoglobulin for GVHD prophylaxis. Several convergent lines of evidence have suggested that inflammatory cytokines act as mediators of acute GVHD.<sup>84</sup> Inflammatory cytokines are implicated in pathophysiology of MF and cause debilitating symptoms as well as mediate higher mortality.<sup>83</sup> Cytokines may also influence the proliferative advantage of neoplastic clone.<sup>82</sup> Could patients with MF be at additional risk of GVHD because of high levels of inflammatory cytokines?

### Poor PS

Symptomatic splenomegaly, debilitating constitutional symptoms, and anemia commonly impair the performance status (PS) of patients with MF. Poor PS at HCT is an independent predictor of a higher NRM and poor survival in MF patients.<sup>56</sup>

## How will the availability of JAK1/2 inhibitors impact the application of HCT in MF?

Although JAK1/2 inhibitor therapy is of significant clinical value in patients with MF, their use neither is curative nor decreases the risk of LT. The impact of the wider availability of JAK1/2 inhibitor therapy on the referral pattern for HCT is not clear. It is possible that some patients who are responding well to JAK1/2 inhibitor therapy may be delayed from consideration for transplantation and may have more advanced-stage disease at the time of actual referral. We think that patients whose therapeutic goal is cure should still be referred for a transplantation consultation, even when they are responding to JAK1/2 inhibitor therapy. This approach is not contradictory because there is a sound theoretical rationale for combining JAK1/2 inhibitor therapy with the transplant conditioning regimen.

**Table 5. Authors' perspective on optimal modern management of MF**

1. MF is a rare chronic hematologic malignancy. It is imperative that patients be referred for a consultation to a center with expertise in the management of MF. The authors believe in a shared care model where the community hematologist/oncologist and the MPN specialist jointly manage the patient with MF.
2. Risk stratification is vital for choosing an optimal treatment strategy and should be done at the time of diagnosis and reviewed periodically during follow-up to identify a change in risk profile.
3. The goals of therapy for each patient need to be defined upfront, taking into consideration factors, such as age, symptom burden, predicted risk of mortality and leukemic transformation, availability of donors, and personal risk tolerance.
4. Therapy should be individualized and include a spectrum of choices ranging from watchful waiting to drug therapy or HCT.
5. Ideally, transplant-eligible patients should make the decision about the choice of initial therapy after receiving a transplantation consultation, and full understanding of risks/benefits associated with HCT versus nontransplantation therapy.
6. Patients experiencing symptom relief and improved performance status on JAK1/2 inhibitor therapy could also consider the option of subsequent HCT.
7. Transplant-eligible patients pursuing a curative goal in the setting of higher-risk MF should proceed to HCT sooner than later, as worsening disease risk score predicts lower posttransplantation survival.
8. Suggested algorithms (Figure 2A-B) and aids to decision making (Figure 3) may further assist in refining treatment decisions in an individual patient.

### Exploring the benefits of combining JAK1/2 inhibitor therapy with HCT


JAK1/2 inhibitor therapy presents an opportunity to address some of the barriers for the success of HCT in MF patients mentioned as discovered in the previous section. JAK1/2 inhibitor therapy is effective in decreasing the burden of troublesome symptoms in patients with MF by reduction of splenomegaly, amelioration of constitutional symptoms, and improvement in PS and well-being.<sup>14,15,50-52</sup> Reduction in splenomegaly may help in faster hematologic recovery in the posttransplantation period. Drugs, such as ruxolitinib, are effective in rapid down-regulation of inflammatory cytokine levels because of anti-JAK1-mediated effect, resulting in improvement of constitutional symptoms.<sup>52</sup> Down-regulation of cytokines may potentially have beneficial impact on GF and acute GVHD.

Improvement in pretransplantation PS and the possible beneficial impact on GF and GVHD make these agents attractive agents for clinical trials in transplant-related strategies for MF. Potential harmful effects of JAK1/2 inhibitor therapy in the transplantation setting may include negative impact on hematologic recovery or explosive splenomegaly or cytokine excesses on withdrawal. Therefore, it is important that the strategy of using JAK1/2 inhibitor therapy in combination with transplantation be explored in a well-designed clinical trial setting wherever possible.

Optimal timing for JAK inhibition in the context of a transplantation strategy appears to be in the pretransplantation setting. In contrast to BCR-ABL inhibitors, which are commonly used to eradicate minimal residual disease in the posttransplantation setting for Philadelphia-positive leukemias, current JAK1/2 inhibitors are not effective in reversing histologic abnormalities or reducing the JAK2V617F allele burden in MF and, therefore, may have limited value in the posttransplantation setting.

### Selection of therapeutic options for MF patients in the JAK inhibitor era

MF is a rare chronic hematologic malignancy. Optimal management of MF will involve close collaboration between community



**Figure 2.** Suggested algorithm for approach to selection of first-line therapy (drug therapy vs HCT). (A) Low/Intermediate-1 risk MF. (B) Intermediate-2/High risk MF.

hematologist/oncologist and a center with expertise in MPN. The author’s perspectives on optimal modern management of MF are summarized in Table 5. In patients, where the goal of therapy is curative, the option of HCT should be considered and a donor search initiated. As proposed by the European LeukemiaNet group, HCT should be considered in patients whose anticipated survival is less than 5 years (usually Intermediate-2/high-risk disease).<sup>85</sup> In addition, transfusion dependency or risk of LT more than or equal to 35% at 3 years would be reasonable indications for considerations for HCT in lower-risk patients. Candidates not eligible for HCT should be offered JAK1/2 inhibitor therapy or clinical trials when symptomatic (Figure 2A-B). Within the framework of this approach, the decision about candidacy for HCT should be

evaluated after careful consideration of the risk posed by disease itself versus risk from transplantation taking into consideration patient-, disease-, and transplant-related factors (Figure 3). Continued study of novel therapeutic strategies, including HCT, is required to optimize patient outcomes in MF.

### Acknowledgments

The authors thank CIBMTR for the data on trends in transplantation for MF as well as Dr Hans Messner (Princess Margaret Hospital) for review of this manuscript and helpful suggestions.




Figure 3. Aids to decision making in selection of initial therapy (drug therapy vs HCT) in patients with MF.

## Authorship

Contribution: V.G. prepared the initial draft of the manuscript; P.H. and R.H. provided further knowledge, insights, discussions, and helped in critical review; and all authors approved the final version of the manuscript.

Conflict-of-interest disclosure: V.G. received clinical trial research funding from Incyte, Novartis, Celgene, YM Biosciences, and Sanofi-Aventis; served on advisory board of Incyte, Novartis,

YM Biosciences, and Sanofi-Aventis; and received consulting fees from Novartis, YM Biosciences, and Sanofi-Aventis. R.H. received research support from Novartis, Celgene, Incyte, Genzyme, AstraZeneca, Bristol Meyers Squibb, and Roche/Genentech. P.H. declares no competing financial interests.

Correspondence: Vikas Gupta, Blood and Marrow Transplant Program, Princess Margaret Hospital, Suite 5-217, 610 University Ave, Toronto, ON, Canada, M5G 2M9; e-mail: vikas.gupta@uhn.ca.

## References

- Tefferi A. Myelofibrosis with myeloid metaplasia. *N Engl J Med*. 2000;342(17):1255-1265.
- Mesa RA, Silverstein MN, Jacobsen SJ, et al. Population-based incidence and survival figures in essential thrombocythemia and agnogenic myeloid metaplasia: an Olmsted County Study, 1976-1995. *Am J Hematol*. 1999;61(1):10-15.
- Tefferi A, Lasho TL, Jimma T, et al. One thousand patients with primary myelofibrosis: the Mayo Clinic experience. *Mayo Clin Proc*. 2012;87(1):25-33.
- Cervantes F, Dupriez B, Pereira A, et al. New prognostic scoring system for primary myelofibrosis based on a study of the International Working Group for Myelofibrosis Research and Treatment. *Blood*. 2009;113(13):2895-2901.
- Gangat N, Caramazza D, Vaidya R, et al. DIPSS plus: a refined Dynamic International Prognostic Scoring System for primary myelofibrosis that incorporates prognostic information from karyotype, platelet count, and transfusion status. *J Clin Oncol*. 2011;29(4):392-397.
- Passamonti F, Cervantes F, Vannucchi AM, et al. A dynamic prognostic model to predict survival in primary myelofibrosis: a study by the IWG-MRT (International Working Group for Myeloproliferative Neoplasms Research and Treatment). *Blood*. 2010;115(9):1703-1708.
- Deeg HJ, Appelbaum FR. Indications for and current results with allogeneic hematopoietic cell transplantation in patients with myelofibrosis. *Blood*. 2011;117(26):7185.
- Tefferi A. How I treat myelofibrosis. *Blood*. 2011;117(13):3494-3504.
- Daly A, Song K, Nevill T, et al. Stem cell transplantation for myelofibrosis: a report from two Canadian centers. *Bone Marrow Transplant*. 2003;32(1):35-40.
- Deeg HJ, Gooley TA, Flowers ME, et al. Allogeneic hematopoietic stem cell transplantation for myelofibrosis. *Blood*. 2003;102(12):3912-3918.
- Guardiola P, Anderson JE, Bandini G, et al. Allogeneic stem cell transplantation for agnogenic myeloid metaplasia: a European Group for Blood and Marrow Transplantation, Societe Francaise de Greffe de Moelle, Gruppo Italiano per il Trapianto del Midollo Osseo, and Fred Hutchinson Cancer Research Center Collaborative Study. *Blood*. 1999;93(9):2831-2838.
- Horan JT, Logan BR, Agovi-Johnson MA, et al. Reducing the risk for transplantation-related mortality after allogeneic hematopoietic cell transplantation: how much progress has been made? *J Clin Oncol*. 2011;29(7):805-813.
- Samuelson S, Sandmaier BM, Heslop HE, et al. Allogeneic haematopoietic cell transplantation for myelofibrosis in 30 patients 60-78 years of age. *Br J Haematol*. 2011;153(1):76-82.
- Verstovsek S, Mesa RA, Gotlib J, et al. A double-blind, placebo-controlled trial of ruxolitinib for myelofibrosis. *N Engl J Med*. 2012;366(9):799-807.
- Harrison C, Kiladjian JJ, Al-Ali HK, et al. JAK inhibition with ruxolitinib versus best available therapy for myelofibrosis. *N Engl J Med*. 2012;366(9):787-798.
- Verstovsek S, Kantarjian HM, Estrov Z, et al. Comparison of outcomes of advanced myelofibrosis patients treated with ruxolitinib (INC018424) to those of a historical control group: survival advantage of ruxolitinib therapy [abstract]. *Blood (ASH Annual Meeting Abstracts)*. 2011;118:793.
- Tefferi A, Litzow MR, Pardanani A. Long-term outcome of treatment with ruxolitinib in myelofibrosis. *N Engl J Med*. 2011;365(15):1455-1457.
- Tefferi A. JAK inhibitors for myeloproliferative neoplasms: clarifying facts from myths. *Blood*. 2012;119(12):2721-2730.
- Dupriez B, Morel P, Demory JL, et al. Prognostic factors in agnogenic myeloid metaplasia: a report on 195 cases with a new scoring system. *Blood*. 1996;88(3):1013-1018.
- Mascarenhas J, Navada S, Malone A, et al. Therapeutic options for patients with myelofibrosis in blast phase. *Leuk Res*. 2010;34(9):1246-1249.
- Mesa RA, Li CY, Ketterling RP, et al. Leukemic transformation in myelofibrosis with myeloid metaplasia: a single-institution experience with 91 cases. *Blood*. 2005;105(3):973-977.
- Tam CS, Nussenzweig RM, Popat U, et al. The natural history and treatment outcome of blast phase BCR-ABL- myeloproliferative neoplasms. *Blood*. 2008;112(5):1628-1637.
- Passamonti F, Cervantes F, Vannucchi AM, et al. Dynamic International Prognostic Scoring System (DIPSS) predicts progression to acute myeloid

- leukemia in primary myelofibrosis. *Blood*. 2010; 116(15):2857-2858.
24. Passamonti F, Rumi E, Elena C, et al. Incidence of leukaemia in patients with primary myelofibrosis and RBC-transfusion-dependence. *Br J Haematol*. 2010;150(6):719-721.
  25. Tefferi A, Pardanani A, Gangat N, et al. Leukemia risk models in primary myelofibrosis: an International Working Group study. *Leukemia*. 2012; 26(6):1439-1441.
  26. Dingli D, Schwager SM, Mesa RA, et al. Presence of unfavorable cytogenetic abnormalities is the strongest predictor of poor survival in secondary myelofibrosis. *Cancer*. 2006;106(9):1985-1989.
  27. Passamonti F, Rumi E, Caramella M, et al. A dynamic prognostic model to predict survival in post-polycythemia vera myelofibrosis. *Blood*. 2008;111(7):3383-3387.
  28. Demory JL, Dupriez B, Fenaux P, et al. Cytogenetic studies and their prognostic significance in agnogenic myeloid metaplasia: a report on 47 cases. *Blood*. 1988;72(3):855-859.
  29. Reilly JT, Snowden JA, Spearing RL, et al. Cytogenetic abnormalities and their prognostic significance in idiopathic myelofibrosis: a study of 106 cases. *Br J Haematol*. 1997;98(1):96-102.
  30. Vaidya R, Caramazza D, Begna KH, et al. Monosomal karyotype in primary myelofibrosis is detrimental to both overall and leukemia-free survival. *Blood*. 2011;117(21):5612-5615.
  31. Hidaka T, Shide K, Shimoda H, et al. The impact of cytogenetic abnormalities on the prognosis of primary myelofibrosis: a prospective survey of 202 cases in Japan. *Eur J Haematol*. 2009;83(4):328-333.
  32. Caramazza D, Begna KH, Gangat N, et al. Refined cytogenetic-risk categorization for overall and leukemia-free survival in primary myelofibrosis: a single center study of 433 patients. *Leukemia*. 2011;25(1):82-88.
  33. Tam CS, Abruzzo LV, Lin KI, et al. The role of cytogenetic abnormalities as a prognostic marker in primary myelofibrosis: applicability at the time of diagnosis and later during disease course. *Blood*. 2009;113(18):4171-4178.
  34. Baxter EJ, Scott LM, Campbell PJ, et al. Acquired mutation of the tyrosine kinase JAK2 in human myeloproliferative disorders. *Lancet*. 2005; 365(9464):1054-1061.
  35. Levine RL, Wadleigh M, Cools J, et al. Activating mutation in the tyrosine kinase JAK2 in polycythemia vera, essential thrombocythemia, and myeloid metaplasia with myelofibrosis. *Cancer Cell*. 2005;7(4):387-397.
  36. Kralovics R, Passamonti F, Buser AS, et al. A gain-of-function mutation of JAK2 in myeloproliferative disorders. *N Engl J Med*. 2005;352(17):1779-1790.
  37. James C, Ugo V, Le Couedic JP, et al. A unique clonal JAK2 mutation leading to constitutive signalling causes polycythaemia vera. *Nature*. 2005; 434(7037):1144-1148.
  38. Scott LM, Tong W, Levine RL, et al. JAK2 exon 12 mutations in polycythemia vera and idiopathic erythrocytosis. *N Engl J Med*. 2007;356(5):459-468.
  39. Oh ST, Simonds EF, Jones C, et al. Novel mutations in the inhibitory adaptor protein LNK drive JAK-STAT signaling in patients with myeloproliferative neoplasms. *Blood*. 2010;116(6):988-992.
  40. Pikman Y, Lee BH, Mercher T, et al. MPLW515L is a novel somatic activating mutation in myelofibrosis with myeloid metaplasia. *PLoS Med*. 2006; 3(7):e270.
  41. Campbell PJ, Griesshammer M, Dohner K, et al. V617F mutation in JAK2 is associated with poorer survival in idiopathic myelofibrosis. *Blood*. 2006;107(5):2098-2100.
  42. Guglielmelli P, Barosi G, Specchia G, et al. Identification of patients with poorer survival in primary myelofibrosis based on the burden of JAK2V617F mutated allele. *Blood*. 2009;114(8):1477-1483.
  43. Tefferi A, Lasho TL, Huang J, et al. Low JAK2V617F allele burden in primary myelofibrosis, compared to either a higher allele burden or unmutated status, is associated with inferior overall and leukemia-free survival. *Leukemia*. 2008; 22(4):756-761.
  44. Barosi G, Bergamaschi G, Marchetti M, et al. JAK2 V617F mutational status predicts progression to large splenomegaly and leukemic transformation in primary myelofibrosis. *Blood*. 2007; 110(12):4030-4036.
  45. Pardanani A, Guglielmelli P, Lasho TL, et al. Primary myelofibrosis with or without mutant MPL: comparison of survival and clinical features involving 603 patients. *Leukemia*. 2011;25(12):1834-1839.
  46. Abdel-Wahab O, Manshouri T, Patel J, et al. Genetic analysis of transforming events that convert chronic myeloproliferative neoplasms to leukemias. *Cancer Res*. 2010;70(2):447-452.
  47. Zhang SJ, Rampal R, Manshouri T, et al. Genetic analysis of patients with leukemic transformation of myeloproliferative neoplasms reveals recurrent SRSF2 mutations which are associated with adverse outcome. *Blood*. 2012;119(19):4480-4485.
  48. Martínez-Trillos A, Gaya A, Maffioli M, et al. Efficacy and tolerability of hydroxyurea in the treatment of the hyperproliferative manifestations of myelofibrosis: results in 40 patients. *Ann Hematol*. 2010;89(12):1233-1237.
  49. Mesa RA, Nagorney DS, Schwager S, et al. Palliative goals, patient selection, and perioperative platelet management: outcomes and lessons from 3 decades of splenectomy for myelofibrosis with myeloid metaplasia at the Mayo Clinic. *Cancer*. 2006;107(2):361-370.
  50. Pardanani A, Gotlib J, Gupta V, et al. An expanded multicenter phase I/II study of CYT387, a JAK-1/2 inhibitor for the treatment of myelofibrosis [abstract]. *Blood (ASH Annual Meeting Abstracts)*. 2011;118:3849.
  51. Pardanani A, Gotlib JR, Jamieson C, et al. Safety and efficacy of TG101348, a selective JAK2 inhibitor, in myelofibrosis. *J Clin Oncol*. 2011;29(7):789-796.
  52. Verstovsek S, Kantarjian H, Mesa RA, et al. Safety and efficacy of INCB018424, a JAK1 and JAK2 inhibitor, in myelofibrosis. *N Engl J Med*. 2010;363(12):1117-1127.
  53. Tyner JW, Bumm TG, Deininger J, et al. CYT387, a novel JAK2 inhibitor, induces hematologic responses and normalizes inflammatory cytokines in murine myeloproliferative neoplasms. *Blood*. 2010;115(25):5232-5240.
  54. Kröger N, Holler E, Kobbe G, et al. Allogeneic stem cell transplantation after reduced-intensity conditioning in patients with myelofibrosis: a prospective, multicenter study of the Chronic Leukemia Working Party of the European Group for Blood and Marrow Transplantation. *Blood*. 2009; 114(26):5264-5270.
  55. Rondelli D, Goldberg JD, Marchioli R, et al. Results of phase II clinical trial MPD-RC 101: allogeneic hematopoietic stem cell transplantation conditioned with fludarabine/melphalan in patients with myelofibrosis [abstract]. *Blood (ASH Annual Meeting Abstracts)*. 2011;118:1750.
  56. Ballen KK, Shrestha S, Sobocinski KA, et al. Outcome of transplantation for myelofibrosis. *Biol Blood Marrow Transplant*. 2010;16(3):358-367.
  57. Scott BL, Gooley TA, Sorror ML, et al. The Dynamic International Prognostic Scoring System for myelofibrosis predicts outcomes after hematopoietic cell transplantation. *Blood*. 2012;119(11):2657-2664.
  58. Kerbauy DM, Gooley TA, Sale GE, et al. Hematopoietic cell transplantation as curative therapy for idiopathic myelofibrosis, advanced polycythemia vera, and essential thrombocythemia. *Biol Blood Marrow Transplant*. 2007;13(3):355-365.
  59. Cervantes F, Rovira M, Urbano-Ispizua A, et al. Complete remission of idiopathic myelofibrosis following donor lymphocyte infusion after failure of allogeneic transplantation: demonstration of a graft-versus-myelofibrosis effect. *Bone Marrow Transplant*. 2000;26(6):697-699.
  60. Devine SM, Hoffman R, Verma A, et al. Allogeneic blood cell transplantation following reduced-intensity conditioning is effective therapy for older patients with myelofibrosis with myeloid metaplasia. *Blood*. 2002;99(6):2255-2258.
  61. Byrne JL, Beshti H, Clark D, et al. Induction of remission after donor leucocyte infusion for the treatment of relapsed chronic idiopathic myelofibrosis following allogeneic transplantation: evidence for a 'graft vs myelofibrosis' effect. *Br J Haematol*. 2000;108(2):430-433.
  62. Bacigalupo A, Ballen K, Rizzo D, et al. Defining the intensity of conditioning regimens: working definitions. *Biol Blood Marrow Transplant*. 2009; 15(12):1628-1633.
  63. Bacigalupo A, Soraru M, Dominietto A, et al. Allogeneic hemopoietic SCT for patients with primary myelofibrosis: a predictive transplant score based on transfusion requirement, spleen size and donor type. *Bone Marrow Transplant*. 2010;45(3):458-463.
  64. Gupta V, Kroger N, Aschan J, et al. A retrospective comparison of conventional intensity conditioning and reduced-intensity conditioning for allogeneic hematopoietic cell transplantation in myelofibrosis. *Bone Marrow Transplant*. 2009; 44(5):317-320.
  65. Patriarca F, Bacigalupo A, Sperotto A, et al. Allogeneic hematopoietic stem cell transplantation in myelofibrosis: the 20-year experience of the Gruppo Italiano Trapianto di Midollo Osseo (GITMO). *Haematologica*. 2008;93(10):1514-1522.
  66. Rondelli D, Barosi G, Bacigalupo A, et al. Allogeneic hematopoietic stem-cell transplantation with reduced-intensity conditioning in intermediate- or high-risk patients with myelofibrosis with myeloid metaplasia. *Blood*. 2005;105(10):4115-4119.
  67. Abellsson J, Merup M, Birgegard G, et al. The outcome of allo-HSCT for 92 patients with myelofibrosis in the Nordic countries. *Bone Marrow Transplant*. 2012;47(3):380-386.
  68. Alchalby H, Yunus DR, Zabelina T, et al. Risk models predicting survival after reduced-intensity transplantation for myelofibrosis [published online ahead of print January 27, 2012]. *Br J Haematol*. doi: 10.1111/j.1365-2141.2011.09009.x.
  69. Nivison-Smith I, Dodds AJ, Butler J, et al. Allogeneic hematopoietic cell transplantation for chronic myelofibrosis in Australia and New Zealand: older recipients receiving myeloablative conditioning at increased mortality risk. *Biol Blood Marrow Transplant*. 2012;18(2):302-308.
  70. Ditschkowski M, Elmaagacli AH, Trensche R, et al. DIPSS scores, pre-transplant therapy and chronic GVHD determine outcome after allogeneic hematopoietic stem cell transplantation for myelofibrosis [published online ahead of print April 4, 2012]. *Haematologica*. doi: 10.3324/haematol.2011.061168.
  71. Alchalby H, Badbaran A, Zabelina T, et al. Impact of JAK2V617F mutation status, allele burden, and clearance after allogeneic stem cell transplantation for myelofibrosis. *Blood*. 2010;116(18):3572-3581.
  72. Ditschkowski M, Elmaagacli AH, Trensche R, et al. No influence of V617F mutation in JAK2 on outcome after allogeneic hematopoietic stem cell transplantation (HSCT) for myelofibrosis. *Biol Blood Marrow Transplant*. 2006;12(12):1350-1351.
  73. Alchalby H, Lioznov M, Fritzsche-Friedland U, et al. Circulating CD34(+) cells as prognostic and

- follow-up marker in patients with myelofibrosis undergoing allo-SCT. *Bone Marrow Transplant.* 2012;47(1):143-145.
74. Takagi S, Ota Y, Uchida N, et al. Successful engraftment after reduced-intensity umbilical cord blood transplantation for myelofibrosis. *Blood.* 2010;116(4):649-652.
75. Li Z, Gooley T, Applebaum FR, et al. Splenectomy and hemopoietic stem cell transplantation for myelofibrosis. *Blood.* 2001;97(7):2180-2181.
76. Stewart WA, Pearce R, Kirkland KE, et al. The role of allogeneic SCT in primary myelofibrosis: a British Society for Blood and Marrow Transplantation study. *Bone Marrow Transplant.* 2010;45(11):1587-1593.
77. Robin M, Esperou H, de Latour RP, et al. Splenectomy after allogeneic haematopoietic stem cell transplantation in patients with primary myelofibrosis. *Br J Haematol.* 2010;150(6):721-724.
78. Kröger N, Alchalby H, Klyuchnikov E, et al. JAK2-V617F-triggered preemptive and salvage adoptive immunotherapy with donor-lymphocyte infusion in patients with myelofibrosis after allogeneic stem cell transplantation. *Blood.* 2009;113(8):1866-1868.
79. Wong KM, Atenafu EG, Kim D, et al. Incidence, risk factors for early hepatotoxicity and its impact on survival in patients with myelofibrosis undergoing allogeneic hematopoietic cell transplantation [published online ahead of print April 21, 2012]. *Biol Blood Marrow Transplant.* doi: 10.1016/j.bbmt.2012.04.011.
80. Bryder D, Ramsfjell V, Dybedal I, et al. Self-renewal of multipotent long-term repopulating hematopoietic stem cells is negatively regulated by Fas and tumor necrosis factor receptor activation. *J Exp Med.* 2001;194(7):941-952.
81. Dybedal I, Bryder D, Fossum A, et al. Tumor necrosis factor (TNF)-mediated activation of the p55 TNF receptor negatively regulates maintenance of cycling reconstituting human hematopoietic stem cells. *Blood.* 2001;98(6):1782-1791.
82. Fleischman AG, Aichberger KJ, Luty SB, et al. TNF $\alpha$  facilitates clonal expansion of JAK2V617F positive cells in myeloproliferative neoplasms. *Blood.* 2011;118(24):6392-6398.
83. Tefferi A, Vaidya R, Caramazza D, et al. Circulating interleukin (IL)-8, IL-2R, IL-12, and IL-15 levels are independently prognostic in primary myelofibrosis: a comprehensive cytokine profiling study. *J Clin Oncol.* 2011;29(10):1356-1363.
84. Ferrara JL, Levine JE, Reddy P, et al. Graft-versus-host disease. *Lancet.* 2009;373(9674):1550-1561.
85. Barbui T, Barosi G, Birgegard G, et al. Philadelphia-negative classical myeloproliferative neoplasms: critical concepts and management recommendations from European LeukemiaNet. *J Clin Oncol.* 2011;29(6):761-770.
86. Tefferi A, Pardanani A, Lim KH, et al. TET2 mutations and their clinical correlates in polycythemia vera, essential thrombocythemia and myelofibrosis. *Leukemia.* 2009;23(5):905-911.
87. Abdel-Wahab O, Pardanani A, Rampal R, et al. DNMT3A mutational analysis in primary myelofibrosis, chronic myelomonocytic leukemia and advanced phases of myeloproliferative neoplasms. *Leukemia.* 2011;25(7):1219-1220.
88. Tefferi A, Jimma T, Sulai NH, et al. IDH mutations in primary myelofibrosis predict leukemic transformation and shortened survival: clinical evidence for leukemogenic collaboration with JAK2V617F. *Leukemia.* 2012;26(3):475-480.
89. Guglielmelli P, Biamonte F, Score J, et al. EZH2 mutational status predicts poor survival in myelofibrosis. *Blood.* 2011;118(19):5227-5234.
90. Score J, Hidalgo-Curtis C, Jones AV, et al. Inactivation of polycomb repressive complex 2 components in myeloproliferative and myelodysplastic/myeloproliferative neoplasms. *Blood.* 2012; 119(5):1208-1213.
91. Puda A, Milosevic JD, Berg T, et al. Frequent deletions of JARID2 in leukemic transformation of chronic myeloid malignancies. *Am J Hematol.* 2012;87(3):245-250.
92. Robin M, Tabrizi R, Mohty M, et al. Allogeneic haematopoietic stem cell transplantation for myelofibrosis: a report of the Societe Francaise de Greffe de Moelle et de Therapie Cellulaire (SFGM-TC). *Br J Haematol.* 2011;152(3):331-339.